

# COSMOLOGICAL CONSTRAINTS FROM THE SPT POL CMB LENSING POWER SPECTRUM

FEDERICO BIANCHINI

w/ KIMMY WU & THE SPT COLLABORATION



THE UNIVERSITY OF  
MELBOURNE

Based on  
[astro-ph:1905.05777](#)  
[astro-ph:1910.07157](#)

# The SPT Collaboration (Aug 2019)

~100 scientists (~half postdocs and students)  
across ~20 institutions




Funded By:




# Outline

- ▶ CMB lensing primer
- ▶ SPTpol: data & instrument
- ▶ SPTpol lensing: measurement & interpretation
- ▶ Looking ahead: SPT-3G



# CMB lensing in a nutshell

CMB photons are weakly gravitationally deflected by the intervening matter distribution during their cosmic journey




$$\tilde{X}(\hat{\mathbf{n}}) = X(\hat{\mathbf{n}} + \nabla\phi(\hat{\mathbf{n}}))$$

$$\phi(\hat{\mathbf{n}}) = -2 \int_0^{\chi_*} d\chi \frac{f_K(\chi_* - \chi)}{f_K(\chi_*) f_K(\chi)} \Psi(\chi \hat{\mathbf{n}}; \eta_0 - \chi)$$


Geometry

Growth of structures



**UNLENSED\***

\*no primordial B-modes



# CMB lensing reconstruction

Lensing introduces **statistical anisotropy**, i.e. correlates previously uncorrelated multipoles

$$\langle X(\mathbf{l})Y^*(\mathbf{l} - \mathbf{L}) \rangle = 0 \xrightarrow{\text{w/ lensing}} \langle X(\mathbf{l})Y^*(\mathbf{l} - \mathbf{L}) \rangle \propto \phi(\mathbf{L})$$

METHOD

We can extract lensing by looking at the off-diagonal correlations between X and Y

$$\bar{\phi}_{\mathbf{L}}^{XY} = \frac{1}{\mathcal{R}_{\mathbf{L}}^{XY}} \int d^2\ell W_{\ell, \ell - \mathbf{L}}^{XY} \bar{X}_\ell \bar{Y}_{\ell - \mathbf{L}}^*$$

Hu&Okamoto02  
Seljak&Zaldarriaga97



Lensing Potential

Normalization



Optimally-chosen  
weight function

Filtered (data) maps  
 $X, Y \in [T, E, B]$


# (a Few) Applications of CMB lensing


**X**



- ▶ Neutrino masses
- ▶ Dark energy
- ▶ Curvature


**X**





- ▶ Tracers bias
- ▶ Shear bias calibration
- ▶ Photo-z uncertainties



B-mode lensing template




- ▶ Delens B-modes
- ▶ Inflationary gravitational waves
- ▶ Scale energy of inflation



# (a Few) Applications of CMB lensing




**X**



- ▶ Neutrino masses

## This talk


- ▶ Curvature




**X**






- ▶ Tracers bias
- ▶ Shear bias calibration
- ▶ Photo-z uncertainties



B-mode lensing template



- ▶ Delens B-modes
- ▶ Inflationary gravitational waves
- ▶ Scale energy of inflation





THE UNIVERSITY OF  
MELBOURNE


# The South Pole Telescope (SPT)



10-meter telescope  
arcmin resolution


## 2007: SPT-SZ

960 detectors  
95,150,220 GHz




## 2012: SPTpol

1600 detectors  
95,150 GHz  
*+polarization*




## 2017: SPT-3G

~16,200 detectors  
95,150, 220 GHz  
*+polarization*




# SPTpol survey



Survey	Obs. Years	Area (deg <sup>2</sup> )	95 GHz (uK- <sup>1</sup> )	150 (uK- <sup>1</sup> )	220 (uK- <sup>1</sup> )
SPT-SZ	2007-11	2500	40	17	80
<b>SPTpol-Main</b>	<b>2012-16</b>	<b>500</b>	<b>12</b>	<b>5</b>	-
SPTpol-Deep	2012-16	100	10	3.5	-

# SPTpol maps


(high-pass filtered) **temperature** map at **150 GHz** for all data  
from 2013 to July 2015



# 500d CMB lensing map

**High S/N map of the projected mass distribution out to the surface of last scattering**

$$\kappa = -\frac{1}{2} \nabla^2 \phi$$





# 500d CMB lensing power spectrum

- ▶ No lensing hypothesis ruled out at  $40\sigma$
- ▶  $S/N > 1$  (sample variance limited) for modes  $L \lesssim 250$

$$A_{\text{MV}} = 0.94 \pm 0.06(\text{stat}) \pm 0.025(\text{sys})$$


$$A_{\text{POL}} = 0.91 \pm 0.09(\text{stat}) \pm 0.04(\text{sys})$$

$$A_{\text{TT}} = 0.83 \pm 0.10(\text{stat}) \pm 0.01(\text{sys})$$



# 500d CMB lensing constraints\*

## $\Lambda$ CDM: $\Omega_m$ vs $\sigma_8$




$$\sigma_8 \Omega_m^{0.25} = 0.593 \pm 0.025 \quad \text{SPTpol CMB lensing only (4\%)}$$

$$\sigma_8 \Omega_m^{0.25} = 0.590 \pm 0.020 \quad \text{Planck CMB lensing only (3\%)}$$

\*“Lensing-only” priors:  $\Omega_b h^2 = 0.0222 \pm 0.0005$ ,  $n_s = 0.96 \pm 0.02$ ,  $0.4 < h < 1$  + fix tau

# 500d CMB lensing constraints


## $\Lambda$ CDM: $\Omega_m$ vs $\sigma_8$



- ▶ Constraining power similar to galaxy lensing  $S_8 = 0.86 \pm 0.11$  (SPTpol lensing)
- ▶ But degeneracy orientation different due to lower-z of source plane

# Adding BAO information

Lensing is sensitive to  $\sigma_8 \Omega_m^{0.25} (\Omega_m h^2)^{-0.37}$ , degeneracies can be broken by including BAO measurements



$$\left. \begin{array}{l} H_0 = 72.0_{-2.5}^{+2.1} \text{ km s}^{-1} \text{ Mpc}^{-1} \\ \sigma_8 = 0.779 \pm 0.023 \\ \Omega_m = 0.368_{-0.037}^{+0.032} \end{array} \right\} \text{SPTPOL lensing} + \text{BAO}$$


$$\left. \begin{array}{l} H_0 = 67.9_{-1.3}^{+1.1} \text{ km s}^{-1} \text{ Mpc}^{-1} \\ \sigma_8 = 0.811 \pm 0.019 \\ \Omega_m = 0.303_{-0.018}^{+0.016} \end{array} \right\} \text{Planck lensing} + \text{BAO}$$

$$\left. \begin{array}{l} H_0 = 72.6_{-2.9}^{+2.3} \text{ km s}^{-1} \text{ Mpc}^{-1} \\ \sigma_8 = 0.814 \pm 0.019 \\ \Omega_m = 0.379_{-0.042}^{+0.036} \end{array} \right\} \text{Planck lensing} \quad \begin{array}{l} 135 < L < 400 \\ + \text{BAO} \end{array}$$

see also Wu+20 (arxiv:2004.10207)



# Joint Constraints from Primary CMB and Lensing

When primary CMB data are included, the lensing power spectrum shape is almost fixed, but the amplitude still has freedom, this helps breaking the  $A_s e^{-2\tau}$  (and geometrical) degeneracy



# Lensing is consistent with $\Lambda$ CDM predictions

Lensing smears the acoustic peaks in the primary CMB TT/TE/EE spectra (effect quantified by the *unphysical* parameter  $A_L$ )




# Massive neutrinos\*

Massive neutrinos suppress the growth of structure below the neutrino free-streaming length, resulting in a scale-dependent suppression of  $C_L^{\phi\phi}$

$\sum m_\nu < 0.20 \text{ eV (95\%)} \text{ (PLANCKTT + LOWP)} + \text{BAO}$

$\sum m_\nu < 0.23 \text{ eV (PLANCKTT + LOWP + BAO} + \text{SPTPOL lensing, 95\%)},$

$\sum m_\nu < 0.22 \text{ eV (PLANCKTT + LOWP + BAO} + \text{Planck lensing, 95\%)},$



\*we consider one single family of massive neutrinos


# Massive neutrinos\*

We can remove the primary CMB sensitivity to lensing by marginalising over  $A_L$

$\sum m_\nu < 0.430 \text{eV} (95\%)$  ( PLANCKTT + LOWP [ $A_L$ ]free)

$\sum m_\nu < 0.45 \text{eV}$  (PLANCKTT + LOWP + BAO  
+ SPTPOL lensing [ $A_L$  free], 95%)


$\sum m_\nu < 0.39 \text{eV}$  (PLANCKTT + LOWP + BAO  
+ Planck lensing [ $A_L$  free], 95%).



\*we consider one single family of massive neutrinos

# Curvature


Primary CMB affected by the geometrical degeneracy: models with similar matter content and angular-diameter distance to the last-scattering surface will produce nearly indistinguishable CMB power spectra...but lensing can break it!



Lensing measurements consistent with a flat LCDM model

# The SPT-3G 1500 deg<sup>2</sup> Survey

*Deep survey complements measurements from a wide survey. For SPT-3G:*





- **Damping Tail Physics:** Improve S/N of CMB power spectra by factors of  $> \sim 10$  at  $\ell > 2500$  over current constraints
- **CMB lensing:** cross-correlations, cluster lensing, de-lensing  $\sigma(r) \sim 0.0025$
- **Clusters:** More efficient at finding high-z clusters, discovering clusters at  $z > 2$ , proto-clusters out to  $z > 4$
- **Transients:** Daily observations of 1500 deg<sup>2</sup> field provide new window into mm-wave transients (e.g., GRBs, FRBs), and mJy-level monitoring of 1000s of blazars
- **and more!**

	Obs. Years	Area (deg <sup>2</sup> )	95 GHz (uK-arcmin)	150 (uK-arcmin)	220 (uK-arcmin)
<b>SPT-3G (projected)</b>	2017-23	1500	3.0	2.2	8.8

# CMB lensing with SPT-3G

Analysis of the 2018 data is underway (led by Zhaodi Pan at UChicago),  
 S/N expected to increase by  $\sim 1.5\times$




Projected uncertainties from full-survey depth

# WRAP-UP

- ▶ **SPTpol provides the most precise CMB Lensing measurement from the ground**
- ▶ **Cosmological constraints are comparable to Planck and consistent with LCDM**
- ▶ **SPT-3G: deep into our 3rd year of observation with 10x more detectors ... stay tuned!**

# THANKS FOR LISTENING!



# Priors

**Table 1**

Summary of the Priors Imposed on Each Cosmological Parameter in This Work, When Considering Either Lensing-only Data Sets or Also Including Primary CMB Measurements

Parameter	Lensing Only	Lensing + CMB
$\Omega_b h^2$	$\mathcal{N}(0.0222, 0.0005^2)$	$\mathcal{U}(0.005, 0.1)$
$\Omega_c h^2$	$\mathcal{U}(0.001, 0.99)$	$\mathcal{U}(0.001, 0.99)$
$H_0$ (km s <sup>-1</sup> Mpc <sup>-1</sup> )	$\mathcal{U}(40, 100)$	$\mathcal{U}(40, 100)$
$\tau$	0.055	$\mathcal{U}(0.01, 0.8)$
$n_s$	$\mathcal{N}(0.96, 0.02^2)$	$\mathcal{U}(0.8, 1.2)$
$\ln(10^{10} A_s)$	$\mathcal{U}(1.61, 3.91)$	$\mathcal{U}(1.61, 3.91)$
$\sum m_\nu$ (eV)	0.06	0.06 or $\mathcal{U}(0, 5)$
$\Omega_K$	0	0 or $\mathcal{U}(-0.3, 0.3)$
$A_L$	1	1 or $\mathcal{U}(0, 10)$
$A_L^{\phi\phi}$	1	1 or $\mathcal{U}(0, 10)$


**Note.** Parameters that are fixed are reported by a single number.  $\mathcal{U}(a, b)$  denotes a uniform distribution between  $[a, b]$ , while  $\mathcal{N}(\mu, \sigma^2)$  indicates a Gaussian distribution with mean  $\mu$  and variance  $\sigma^2$ .

# CMB lensing likelihood correction


Fiducial cosmological parameters  $\theta$  enter the CMB lensing likelihood in several ways:

- Directly through the theory power spectrum  $C_l \Phi\Phi$
- Indirectly (non-linearly) through the CMB power spectra needed to calculate the response functions
- Indirectly (linearly) through  $C_l \Phi\Phi$  that appears in the N1 calculation

$$\begin{aligned}
 C_L^{\phi\phi} &= [C_L^{\phi\phi}|_\theta] + \sum_{XY,WZ} w_L^{XY} w_L^{ZW} \sum_{j \in [TT,TE,EE]} \boxed{\frac{\partial \ln(\mathcal{R}^{XY}\mathcal{R}^{ZW})}{\partial C_{L'}^j}} C_L^{\phi\phi}|_{\text{fid}} (C_{L'}^j|_\theta - C_{L'}^j|_{\text{fid}}) + \\
 &\quad \sum_{XY,WZ} w_L^{XY} w_L^{ZW} \boxed{\frac{\partial N_L^{1,XYZW}}{\partial C_{L'}^{\phi\phi}}} (C_{L'}^{\phi\phi}|_\theta - C_{L'}^{\phi\phi}|_{\text{fid}}) \\
 &= C_L^{\phi\phi}|_\theta + \sum_{j \in [TT,TE,EE,\phi\phi]} M_{LL'}^j (C_{L'}^j|_\theta - C_{L'}^j|_{\text{fid}}) \\
 -2 \ln \mathcal{L}_\phi(\Theta) &= \sum_{ij} [\hat{C}_{L_b^i}^{\phi\phi} - C_{L_b^i}^{\phi\phi,\text{th}}(\Theta)] \mathbb{C}_{L_b^i L_b^j}^{-1} [\hat{C}_{L_b^j}^{\phi\phi} - C_{L_b^j}^{\phi\phi,\text{th}}(\Theta)],
 \end{aligned}$$



# Other CMB lensing power spectra



# More on the BAO + CMB Lensing (+BBN)




FIG. 1. The posterior distributions of the data sets BAO+BBN, BAO+BBN+SPTpol lensing, BAO+BBN+*Planck* lensing in the parameter basis  $[\theta_{\perp}, \theta_{\parallel}, \sigma_8 \Omega_m^{1/4}]$  which is native to the BAO and CMB lensing measurements. The posteriors for the combined BAO+BBN+*Planck* lensing+SPTpol lensing data set are not shown for clarity as they are qualitatively close to those from BAO+BBN+*Planck* lensing.






FIG. 2. The posterior distributions of the data sets BAO+BBN, BAO+BBN+SPTpol lensing, BAO+BBN+*Planck* lensing for  $H_0$  and  $\theta_{\parallel}$ . The differences observed in  $H_0$  are highly correlated with the differences in  $\theta_{\parallel}$ , which suggests a common origin in the parallel BAO measurements.

# Other lensing measurements w/ SPT



## Cluster lensing



Raghunathan ,...,FB,...+ (SPT), 18